

ADULT SEPSIS CRITERIA

SIRS	<ul style="list-style-type: none"> • Temperature >38.3 or < 36.0 C • Heart Rate (pulse) > 90 • Respirations > 20 per minute • WBC >12,000 or < 4,000 or >10% bands
Sepsis	<ul style="list-style-type: none"> • Known or suspected infection PLUS two or more SIRS criteria
Severe Sepsis (Sepsis PLUS New Organ Failure)	<p style="text-align: center; margin: 0;"><small>Organ Failure Criteria</small></p> <ul style="list-style-type: none"> • SBP < 90 or MAP <65 • Acute respiratory failure as evidenced by a new need for invasive (e.g. intubation) or non-invasive (BIPAP) ventilation • Creatinine > 2 • Urine output < 0.5ml/kg/hour for 2 hrs. • Bilirubin > 2 mg/dL • Platelet count < 100,000 • INR > 1.5 • aPTT > 60 sec • Lactate > 2 mmol/L • Altered level of consciousness
Septic Shock	<ul style="list-style-type: none"> • Severe sepsis plus hypotension despite fluid bolus of 30ml/kg • Lactate \geq 4mmol/L

****severe sepsis presentation begins at the moment the last of the clinical criteria for severe sepsis are met or physician documentation of severe sepsis or septic shock ****

Core Measure Sepsis Bundle to be completed within 3 hours of severe sepsis presentation

Print and complete sepsis checklist from standard register-form # NS-624B